

OU-HET 278

hep-ph/9709367

September, 1997

Reconstruction of quark mass matrices in the NNI form from the experimental data

Eiichi Takasugi* and Masaki Yoshimura†

*Department of Physics, Osaka University
1-16 Toyonaka, Osaka 560, Japan*

Abstract

We examined the question that what is a general form of quark mass matrices which is achieved by the transformation that leaves the left-handed gauge interaction invariant. In particular, we analyzed in detail the Fritzsch-type and the Branco-Silva-Marcos-type parametrization. Both parametrizations contain ten parameters and can be expressed by the experimental data. We explicitly reconstructed quark mass matrices in terms of quark masses and CKM parameters for the Fritzsch-type parametrization.

* e-mail address: takasugi@phys.wani.osaka-u.ac.jp

† e-mail address: masaki@phys.wani.osaka-u.ac.jp

1 Introduction

There are increasing interests in understanding the CKM mixings and the CP violation parameter. Many works were made to find *Ansätze* for quark mass matrices and discussed its predictions. The Fritzsch *Ansatz*[1] is one of examples. This approach is called the texture zero analysis where some elements of mass matrices are required to be zero to reduce the degrees of freedom in mass matrices. The extensive works are made along this line[2]-[5].

Recently, some attention was paid to the inverse problem, that is, to reconstruct quark mass matrices directly from the experimental data. This problem is ambiguous until the form of quark mass matrices is fixed. Therefore, we need to choose a general and convenient form of quark mass matrices to start with. One of candidates is the nearest neighbor interaction (NNI) form which is considered as a “general” form of quark mass matrices because this form is achieved by the transformation that leaves the left-handed gauge interaction invariant[6]. Harayama and Okamura[7] chose the NNI form and showed directly how elements of quark mass matrices are expressed by observables, quark masses and CKM parameters. However, the NNI form of quark mass matrices contain twelve parameters so that all parameters are not fixed uniquely because there are only ten observables for three generation case. Hereafter, we consider only three generation case. Subsequently, Koide[8] showed that within the NNI form the 3-2 element of up-quark mass matrix can be made zero in general. This is achieved by using the rephasing freedom of quarks. Now the number of parameters in this parametrization is ten so that quark mass matrices are completely determined by the data.

Recently, one of authors showed[9] that quark mass matrices can be transformed in general to either one of the following two forms, the Fritzsch-type parametrization or the Branco-Silva-Marcos(BS)-type parametrization. The Fritzsch-type parametrization is that the u-quark mass matrix is in a Fritzsch form[1], while the d-quark mass matrix is in the NNI form. The BS-type parametrization is that the d-quark mass matrix is in the Branco-Silva-Marcos form[10], while the u-quark mass matrix is in the NNI form. The

transformation to these forms is constructed explicitly in Ref.[9] and examined for what quark mass matrices the transformation is possible. In both parametrizations, quark mass matrices contain ten physical parameters so that they can be determined uniquely by quark masses and the CKM parameters. The formula how to express quark mass matrices in terms of data are given. Falcone, Pisanti and Rosa[11] reconstructed quark mass matrices by using some special form: M_u is in a diagonal form and M_d has zero at the 1-1, 2-2 and 3-1 elements, and the form where M_u and M_d are exchanged.

In this paper, we analyze the transformation in detail and examine for what form of quark mass matrices the transformation to the Fritzsch-type or the BS-type parametrization is possible. Our answer is that the transformation is possible for most of physically meaningful quark mass matrices, that is, those which reproduce the present data of quark masses and CKM mixing parameters well. Then we reconstruct quark mass matrices from the experimental data in the Fritzsch-type parametrization.

In below, we shall briefly explain the transformation as an introduction. First, we give the transformation of quark mass matrices to the Fritzsch form:

$$U^\dagger M_u V_u = \widetilde{M}_{uF} , \quad U^\dagger M_d V_d = \widetilde{M}_d , \quad (1)$$

where

$$\widetilde{M}_{uF} = \begin{pmatrix} 0 & a_u & 0 \\ a_u & 0 & c_u \\ 0 & c_u & e_u \end{pmatrix} , \quad (2)$$

and

$$\widetilde{M}_d = \begin{pmatrix} 0 & a_d e^{i\alpha_1} & 0 \\ b_d & 0 & c_d e^{i\alpha_2} \\ 0 & d_d & e_d \end{pmatrix} , \quad (3)$$

under the condition that the equation for the u-quark phases θ_j ,

$$\sum_{j,k} e^{i(\theta_j - \theta_k)} (O_u)_{1j} (O_u)_{2k} (K D_d^2 K^\dagger)_{jk} = 0 , \quad (4)$$

has the solution. This equation contains two phases $\theta_{12} \equiv \theta_1 - \theta_2$ and $\theta_{23} \equiv \theta_2 - \theta_3$ which are to be fixed by this equation. Here, D_d is the diagonal mass matrix for d-quarks, K is

the CKM matrix and O_u is the orthogonal matrix which diagonalizes the Fritzsch matrix $\widetilde{M}_{uF}$. Elements of u-quark matrix $\widetilde{M}_{uF}$ are written explicitly by the u-quark masses and d-quark mass matrix $\widetilde{M}_d$ are given by

$$\begin{aligned}
a_d &= \sqrt{H_{d11}} , & b_d &= \sqrt{\frac{\det H_d}{H_{d11}H_{d33} - |H_{d13}|^2}} , \\
c_d &= \frac{|H_{d23}|\sqrt{H_{d11}}}{\sqrt{H_{d11}H_{d33} - |H_{d13}|^2}} , & d_d &= \frac{|H_{d13}|}{\sqrt{H_{d11}}} , \\
e_d &= \frac{\sqrt{H_{d11}H_{d33} - |H_{d13}|^2}}{\sqrt{H_{d11}}} , \\
\alpha_1 &= \arg H_{d13} , & \alpha_2 &= \arg H_{d23} ,
\end{aligned} \tag{5}$$

where elements of the matrix H_d are given by

$$(H_d)_{il} = \sum_{j,k} e^{i(\theta_j - \theta_k)} (O_u)_{ij} (O_u)_{lk} (K D_d^2 K^\dagger)_{jk} . \tag{6}$$

The matrix H_d contains u-quark phases which are determined from Eq.(4) and all others are determined by quark masses and the CKM matrix.

Similarly, the transformation of quark mass matrices to the BS form is obtained. Here, M_d is transformed to the BS form

$$\widetilde{M}_{dBS} = \begin{pmatrix} 0 & a_d & 0 \\ a_d & 0 & c_d \\ 0 & e_d & e_d \end{pmatrix} , \tag{7}$$

while all others are obtained by just changing the suffix u to d and the phase θ_j to ϕ_j and K to $K^\dagger$. Thus the condition for the existence of the transformation becomes

$$\sum_{j,k} e^{i(\phi_j - \phi_k)} (O_d)_{1j} (O_d)_{2k} (K^\dagger D_u^2 K)_{jk} = 0 . \tag{8}$$

Here, D_u is the diagonal mass matrix for u-quarks and O_d is the orthogonal matrix which diagonalizes $\widetilde{M}_{dBS} \widetilde{M}_{dBS}^\dagger$.

In the following, we solve equations in Eq.(4) and (8) and examine in what region of CKM parameters the solution exists and reconstruct the quark mass matrices from the experimental data for the Fritzsch-type parametrization.

In Sec.2, we see how M_u can be transformed to a Fritzsch form or how M_d can be transformed to a BS form. We obtain a complex equation containing two phases for each case. In Sec.3, we show that these equations have solutions for two phases, when mass matrices are ones which reproduce the CKM quark mixings. Summary is given in Sec.4.

2 Region of CKM parameters where the transformation exists

We examine the equation for quark phases given in Eq.(4) or (8). If the solution exists, the transformation of quark mass matrices to the Fritzsch-type form or the BS-type form becomes possible. It is hard to solve these equation explicitly so that we try to solve perturbatively with respect to a small parameter, Cabibbo angle λ . For this purpose, we first parametrize the CKM matrix in a Wolfenstein form

$$K \simeq \begin{pmatrix} 1 - \frac{\lambda^2}{2} & \lambda & \lambda^4 \sigma e^{i\delta} \\ -\lambda & 1 - \frac{\lambda^2}{2} & \lambda^2 A \\ \lambda^3 \sigma' e^{i\delta'} & -\lambda^2 A & 1 \end{pmatrix}, \quad (9)$$

where σ , σ' , δ and δ' are related to ρ and η in the original Wolfenstein parameter as

$$A\rho = \lambda\sigma \cos \delta = A - \sigma' \cos \delta', \quad A\eta = \lambda\sigma \sin \delta = \sigma' \sin \delta'. \quad (10)$$

We define ratios of quark masses as

$$\begin{aligned} r_u &\equiv (m_u/m_c)/\lambda^4 \sim O(1), & r_c &\equiv (m_c/m_t)/\lambda^4 \sim O(1), \\ r_d &\equiv (m_d/m_s)/\lambda^2 \sim O(1), & r_s &\equiv (m_s/m_b)/\lambda^{5/2} \sim O(1). \end{aligned} \quad (11)$$

When we need to estimate numerically, we use $\lambda = 0.2205$ for the Cabibbo angle and the running quark masses in units of GeV defined at $\mu = M_Z$ [12],

$$\begin{aligned} m_u &= 0.00222, & m_c &= 0.661, & m_t &= 180, \\ m_d &= 0.00442, & m_s &= 0.0847, & m_b &= 2.996. \end{aligned} \quad (12)$$

2.1 The case of the Fritzsche-type parametrization

In order to solve the equation (4) for phases, $\theta_{12} = \theta_1 - \theta_2$ and $\theta_{23} = \theta_2 - \theta_3$, we need the orthogonal matrix O_u which diagonalizes the Fritzsche mass matrix for u-quark and $KD_d^2K^\dagger$. The matrix O_u is expressed in terms of ratios of up-quark masses as

$$O_u \simeq \begin{pmatrix} 1 & -\lambda^2\sqrt{r_u} & \lambda^8 r_c \sqrt{r_u r_c} \\ \lambda^2\sqrt{r_u} & 1 & \lambda^2\sqrt{r_c} \\ -\lambda^4\sqrt{r_u r_c} & -\lambda^2\sqrt{r_c} & 1 \end{pmatrix}. \quad (13)$$

Next, the matrix $KD_d^2K^\dagger$ is estimated by using d-quark masses and CKM parameters. In the leading order of λ , we find

$$\begin{aligned} (KD_d^2K^\dagger)_{11} &\simeq m_b^2\lambda^7(r_s^2 + \lambda\sigma^2), & (KD_d^2K^\dagger)_{12} &\simeq m_b^2\lambda^6(A\sigma e^{-i\delta} + r_s^2), \\ (KD_d^2K^\dagger)_{22} &\simeq m_b^2\lambda^4(A^2 + \lambda r_s^2), & (KD_d^2K^\dagger)_{13} &\simeq m_b^2\lambda^4\sigma e^{-i\delta}, \\ (KD_d^2K^\dagger)_{33} &\simeq m_b^2, & (KD_d^2K^\dagger)_{23} &\simeq m_b^2\lambda^2 A. \end{aligned} \quad (14)$$

By keeping leading order terms of λ , the equation (4) is expressed as

$$A_1 e^{i\theta_{12}} - B_1 e^{i\theta_{23}} + C_1 e^{i(\theta_{12} + \theta_{23} - \delta)} = D_1, \quad (15)$$

where

$$\begin{aligned} A_1 &= \lambda^6(A\sigma e^{-i\delta} + r_s^2) \equiv |A_1|e^{-i\kappa}, & B_1 &= \lambda^6 A \sqrt{r_u r_c}, \\ C_1 &= \lambda^6 \sigma \sqrt{r_c}, & D_1 &= \lambda^6 \sqrt{r_u}(A^2 + \lambda r_s^2). \end{aligned} \quad (16)$$

This is a complex valued equation with two variables θ_{12} and θ_{23} . From the equation $|\exp(i(\theta_{12} - \delta)) / (|A_1| \exp(-i(\kappa - \delta)) + C_1 \exp(i\theta_{23}))| = |(D_1 + B_1 e^{i\theta_{23}})|$, we find

$$\cos(\theta_{23} - \zeta) = \frac{|A_1|^2 + C_1^2 - B_1^2 - D_1^2}{2N} \equiv \frac{Z}{2N}, \quad (17)$$

where,

$$\begin{aligned} N &= \sqrt{p^2 + q^2 + 2pq \cos \delta}, \\ \cos \zeta &= -(p + q \cos \delta)/N, \\ \sin \zeta &= -(q \sin \delta)/N, \end{aligned} \quad (18)$$

with

$$\begin{aligned} p &= \lambda^{12} A \sqrt{r_c} [\sigma^2 - r_u (A^2 + \lambda r_s^2)] , \\ q &= \lambda^{12} \sigma r_s^2 \sqrt{r_c} . \end{aligned} \quad (19)$$

The condition of the existence of the solution of θ_{23} and also θ_{12} is $-1 \leq Z/2N \leq 1$. By using quark masses in Eq.(12) and λ , this condition gives the region in the $\rho - \eta$ plain for a given A . In Fig.1 we show this region by taking into account one standard deviation of the experimental value A . In these figures, a region surrounded by the various circles is an experimentally allowed area. The dotted area is the region where the solution exists, that is, the transformation to the Fritzsch-type parametrization is possible. This region is almost independent of the change of A . From these figures, we can say that the transformation is possible for most physically allowed region except for a small area in the first quadrant of CP phase. The vertex point of the triangle corresponds to the Fritzsch-BS *Ansatz* case where a Fritzsch form is assumed for the up-quark mass matrix and the BS form for the d-quark one. This *Ansatz* contains eight free parameters so that the position in the $\rho - \eta$ plane is fixed [13].

2.2 The case of the BS-type parametrization

Analysis can be made similarly to the Fritzsch case. The matrix O_d which diagonalizes the BS mass matrix for d-quarks is given by

$$O_d \simeq \begin{pmatrix} 1 & -2^{-1/4} \lambda \sqrt{r_d} & 2^{-1/4} \lambda^{7/2} r_s \sqrt{r_d} \\ 2^{-1/4} \lambda \sqrt{r_d} & 1 & \lambda^{5/2} r_s \\ -2^{3/4} \lambda^{7/2} r_s \sqrt{r_d} & -\lambda^{5/2} r_s & 1 \end{pmatrix} \quad (20)$$

and $K^\dagger D_u^2 K$ is given by

$$\begin{aligned} (K^\dagger D_u^2 K)_{11} &\simeq m_t^2 \lambda^6 \sigma'^2 , & (K^\dagger D_u^2 K)_{12} &\simeq -m_t^2 \lambda^5 A \sigma' e^{-i\delta'} , \\ (K^\dagger D_u^2 K)_{22} &\simeq m_t^2 \lambda^4 A^2 , & (K^\dagger D_u^2 K)_{13} &\simeq m_t^2 \lambda^3 \sigma' e^{-i\delta'} , \\ (K^\dagger D_u^2 K)_{33} &\simeq m_t^2 , & (K^\dagger D_u^2 K)_{23} &\simeq -m_t^2 \lambda^2 A . \end{aligned} \quad (21)$$

By keeping leading order terms, the equation (8) is expressed as

$$A_2 e^{i(\phi_{12} - \delta')} - i B_2 \sin \phi_{23} - C_2 e^{i(\phi_{12} - \delta' + \phi_{23})} = -D_2 , \quad (22)$$

where $\phi_{12} = \phi_1 - \phi_2$, $\phi_{23} = \phi_2 - \phi_3$, and

$$\begin{aligned} A_2 &= \lambda^5 A \sigma' , & B_2 &= 2^{3/4} \lambda^{11/2} A r_s \sqrt{r_d} , \\ C_2 &= \lambda^{11/2} \sigma' r_s , & D_2 &= 2^{-1/4} \lambda^5 \sqrt{r_d} (A^2 - \lambda r_s^2) . \end{aligned} \quad (23)$$

We find a solution for ϕ_{23} as

$$\cos \phi_{23} = \lambda^{-1/2} \left[\sqrt{2} \sigma'^2 / r_d - (A^2 + \lambda r_s^2) \right] / 2 A r_s . \quad (24)$$

There is another solution, $\lambda^{-1/2} (A^2 + \lambda r_s^2) / 2 A r_s$, but this is unphysical because it is larger than one if we use the current CKM data and quark masses given in Eq.(12). Once ϕ_{23} has a solution, ϕ_{12} also has a solution. We show the result in Fig.2, where the dotted area is an allowed region similarly to the Fritzsch-type parametrization case. This region is almost independent of the change of A and extends all experimentally allowed area. So we conclude that the transformation to the BS-type parametrization is possible for all allowed parameters region. The point indicated by F-BS type corresponds to the case where M_u takes a Fritzsch form and M_d does a BS form where unique prediction for ρ and η is obtained.

3 Reconstruction of the quark mass matrices for the Fritzsch-type parametrization

The u-quark mass matrix is in the Fritzsch form so that it is expressed by u-quark masses so that we concentrate $\widetilde{M}_d$. This can be made by using a formula given in Eq.(5). For this, we have to evaluate H_d first and the angles θ_{12} and θ_{23} next. By using the expressions in (13) and (14), we find

$$\begin{aligned} H_{d11} &\simeq m_b^2 \lambda^7 \left[r_s^2 + \lambda \left\{ \sigma^2 + A^2 r_u - 2 \sqrt{r_u} (A \sigma \cos(\theta_{12} - \delta) + r_s^2 \cos \theta_{12}) \right\} \right] , \\ H_{d22} &\simeq m_b^2 \lambda^4 \left[A^2 + r_c + 2 A \sqrt{r_c} \cos \theta_{23} + \lambda r_s^2 \right] , \\ H_{d33} &\simeq m_b^2 , \\ H_{d13} &\simeq m_b^2 \lambda^4 \left[\sigma e^{i(\theta_{12} - \delta + \theta_{23})} - A \sqrt{r_u} e^{i\theta_{23}} \right] , \\ H_{d23} &\simeq m_b^2 \lambda^2 \left[A e^{i\theta_{23}} + \sqrt{r_c} \right] . \end{aligned} \quad (25)$$

In addition, we have

$$\begin{aligned}\det H_d &= m_d^2 m_s^2 m_b^2 = \lambda^{14} r_d^2 r_s^4 m_b^6, \\ \text{tr} H_d &= m_d^2 + m_s^2 + m_b^2 = (1 + \lambda^5 r_s^2 + \lambda^9 r_s^2 r_d^2) m_b^2.\end{aligned}\tag{26}$$

Then, the elements of $\widetilde{M}_d$ are reconstructed as

$$\begin{aligned}a_d &\simeq m_b \lambda^{7/2} \sqrt{r_s^2 + \lambda f_1}, \\ b_d &\simeq m_b \lambda^{7/2} r_d r_s^2 \sqrt{\frac{1}{r_s^2 - \lambda f_2}}, \\ c_d &\simeq m_b \lambda^2 \sqrt{\frac{(A^2 + r_c + f_3)(r_s^2 + \lambda f_1)}{r_s^2 - \lambda f_2}}, \\ d_d &\simeq m_b \lambda^{1/2} \sqrt{\frac{\sigma^2 + A^2 r_u - f_4}{r_s^2 + \lambda f_1}}, \\ e_d &\simeq m_b \sqrt{\frac{r_s^2 - \lambda f_2}{r_s^2 + \lambda f_1}}, \\ \alpha_1 &\simeq \theta_{23} + \tan^{-1} \left(\frac{\sigma \sin(\theta_{12} - \delta)}{\sigma \cos(\theta_{12} - \delta) - A \sqrt{r_u}} \right), \\ \alpha_2 &\simeq \tan^{-1} \left(\frac{A \sin \theta_{23}}{A \cos \theta_{23} + \sqrt{r_c}} \right)\end{aligned}\tag{27}$$

where

$$\begin{aligned}f_1 &\equiv \sigma^2 + A^2 r_u - 2\sqrt{r_u}(A\sigma \cos(\theta_{12} - \delta) + r_s^2 \cos \theta_{12}), \\ f_2 &\equiv 2\sqrt{r_u} r_s^2 \cos \theta_{12}, \\ f_3 &\equiv 2A\sqrt{r_c} \cos \theta_{23}, \\ f_4 &\equiv 2A\sigma \sqrt{r_u} \cos(\theta_{12} - \delta).\end{aligned}\tag{28}$$

These expressions contain phases θ_{12} , $\theta_{12} - \delta$ and θ_{23} which are fixed by solving Eq.(4) or (15). That is, from Eq.(17), $\cos(\theta_{23} - \zeta)$ is given and then the angle θ_{23} is obtained as

$$\begin{aligned}\cos \theta_{23} &= \left[-Z(p + q \cos \delta) \pm \sqrt{4N^2 - Z^2} q \sin \delta \right] / 2N^2, \\ \sin \theta_{23} &= - \left[\pm \sqrt{4N^2 - Z^2} (p + q \cos \delta) + Zq \sin \delta \right] / 2N^2,\end{aligned}\tag{29}$$

where the upper sign corresponds to positive sign case of $\sin(\theta_{23} - \zeta)$ and the lower sign does to negative sign. Next, by using $\exp(i(\theta_{12} - \delta)) = (D_1 + B_1 e^{i\theta_{23}}) / (|A_1| \exp(-i(\kappa - \delta))) +$

$C_1 \exp(i\theta_{23})$) we obtain

$$\begin{aligned}\cos(\theta_{12} - \delta) &= \left[B_1 C_1 + \lambda^6 D_1 A \sigma + D_1 \lambda^6 r_s^2 \cos \delta + B_1 \lambda^6 r_s^2 \sin \delta \sin \theta_{23} \right. \\ &\quad \left. + (C_1 D_1 + \lambda^6 B_1 A \sigma + B_1 \lambda^6 r_s^2 \cos \delta) \cos \theta_{23} \right] / M , \\ \sin(\theta_{12} - \delta) &= \left[\left\{ \lambda^6 B_1 (A \sigma + r_s^2 \cos \delta) - C_1 D_1 \right\} \sin \theta_{23} \right. \\ &\quad \left. - (D_1 + B_1 \cos \theta_{23}) \lambda^6 r_s^2 \sin \delta \right] / M ,\end{aligned}\tag{30}$$

where,

$$\begin{aligned}M &\equiv (\lambda^6 A \sigma)^2 + (\lambda^6 r_s^2)^2 + C_1^2 + 2 A \sigma \lambda^{12} r_s^2 \cos \delta \\ &\quad + 2 \lambda^6 C_1 \left\{ (r_s^2 \cos \delta + A \sigma) \cos \theta_{23} + r_s^2 \sin \delta \sin \theta_{23} \right\} .\end{aligned}\tag{31}$$

Finally, θ_{12} is obtained by using Eq.(30).

By substituting these angles, the mass matrix M_d is expressed in terms of quark masses and CKM parameters.

Since expressions are complicated, we show the result numerically. For quark masses and λ , we take the values given before and we use the central values for A and σ , i.e., $|K_{cb}|$ and $|K_{ub}|$. Then, elements of $\widetilde{M}_d$ depend on the CP violating parameter δ . Thus, we present the values of elements as a function of δ . We show the result in Fig.3 and Fig.4 which correspond to positive and negative values of $\sin(\theta_{12} - \zeta)$, respectively. The region where the transformation is possible is $\cos \delta \leq 0.66$ [9].

These figures show that there is no symmetric matrices, i.e., $|a_d| = b_d$ and $|c_d| = d_d$ in the NNI form which reproduce the present data. The constraint $|a_d| = b_d$ is possible for $\delta \sim 130^\circ, \sim 60^\circ$, but $|c_d|$ is different from d_d . In Fig.4, we see that $|a_d|$ approximately equals b_d , and d_d does e_d for $\delta \sim 60^\circ$ which is the case where the Fritzsch *Ansatz* for M_u and the BS *Ansatz* for M_d .

4 Discussions

We showed that the transformation of quark mass matrices which leaves the left-handed gauge interaction invariant to the Fritzsch-type form or to the BS-type form is possible in

“general”. The word “general” means that it is possible for almost all quark mass matrices which reproduce the present data of CKM mixings and the CP violating phase. Thus, we concluded that the above forms are most general forms which we can start in general. By using the Fritzsch-type form, we reconstructed quark mass matrices from quark masses, CKM mixings and the CP violating parameter. We calculated each elements of mass matrices numerically and showed in Figs.3 and 4. As it is already known, the symmetric mass matrices are not allowed in this parametrization so that the asymmetric form is unavoidable.

In this paper, we discussed quark mass matrices at M_Z scale, but it is more interesting to have them at the unification scale since they may be related directly to the structure of Yukawa interactions in grand unified theories. This can be done immediately by following our procedure once the data at the unification scale are given. Thus, it is important to obtain the CKM mixings at the unification scale from those at the weak scale, which is now under consideration.

The asymmetric form of quark mass matrices is inevitable for the NNI basis so that we have to go to another type form if we prefer to use symmetric form. The symmetric form arises naturally in models like SO(10) unification. As for the symmetric form, the most interesting choice will be the one where the 1-1 and 1-3 elements are zero. It should be possible to transform quark masses in this form in general and reconstruct the quark mass matrices from the data. This is now under investigation.

Finally, from our analysis we can say what quark mass matrices are required once the CP violating phase is fixed in the Fritzsch-type NNI parametrization.

References

- [1] H. Fritzsch , Phys. Lett. **B73** (1978) 317 ; Nucl. Phys. **B115** (1979) 189.
- [2] S. Dimopoulos , L. J. Hall and S. Raby , Phys. Rev. Lett. **68** (1992) 1984.
- [3] P. Ramond , R. G. Robert and G. G. Ross , Nucl. Phys. **B406** (1993) 19.
- [4] G. Anderson , S. Dimopoulos , L. J. Hall , S. Raby and G. D. Starkman , Phys. Rev. **D49** (1994) 3660.
- [5] K. S. Babu and R. N. Mohapatra , Phys. Rev. Lett. **74** (1995) 2418.
- [6] G. C. Branco , L. Lavoura and F. Mota , Phys. Rev. **D39** (1989) 3443.
- [7] K. Harayama and N. Okamura , Phys. Lett. **B387** (1996) 614.
- [8] Y. Koide , US-97-01 , hep-ph/9701345.
- [9] E. Takasugi , OU-HET 264 , hep-ph/9705263.
- [10] G. C. Branco and J. I. Silva-Marcos , Phys. Lett. **B331** (1994) 390.
- [11] D. Falcone , O. Pisanti and L. Rosa , DSF-T-97/25 , hep-ph/9706300.
- [12] H. Fusaoka and Y. Koide , AMU-97-01 , US-97-03 , hep-ph/9706211.
- [13] T. Ito , Prog. Theor. Phys. **96** (1996) 1055.

Figure caption

- Fig.1 : The CKM parameter region where the transformation to the Fritzsche-type parametrization is possible. The allowed region is shown by the dotted area which is almost independent of A . These figures show that for most of physical quark mass matrices the transformation is possible. The F-BS point shows the predicted point when the Fritzsche form is assumed for M_u and the BS form for M_d .
- Fig.2 : The CKM parameter region where the transformation to the BS-type parametrization is possible. These figures show that for all physical quark mass matrices the transformation is possible.
- Fig.3 : The reconstructed elements of M_d in the Fritzsche-type parametrization for $\sin(\theta_{12} - \zeta) > 0$. In this case, M_u is in the Fritzsche form and is expressed by u-quark masses.
- Fig.4 : The reconstructed elements of M_d in the Fritzsche-type parametrization for $\sin(\theta_{12} - \zeta) < 0$. The F-BS point corresponds to the $\delta \sim 60^\circ$ where $|a_d| = b_d$ and $d_d \simeq e_d$.

Fig.1






Fig.2






Fig.3





Fig.4

